Preparing for your Colonoscopy with SuTab (am appt)

Patient Name:	Physician:
Procedure date:	ARRIVAL TIME:
Location:	Questions: 571.248.0653 option WEEK BEFORE PROCEDURE:
PLEASE purchase the FOLLOWING INSuTab prescription at your	
Peppermint candy (optional, helps t	•
Bottle of MIRALAX (PEG3350) if your property of the prope	•
7 DAYS BEFORE PROCEDURE:	
	ovements a week OR BMI over 35, start taking 1 capful (17 grams) of
Miralax every morning. This is pure	chased separately as indicated above.
 If you are taking Ozempic, Wegovy stop for at least 7 days. 	y, Mounjaro, Rybelius, Trulicity, Victorza or other GLP1 medication,
5 DAYS BEFORE PROCEDURE:	
	mato skins, raw vegetables, corn, popcorn, bran, and chia seeds.
 STOP iron supplements, vitamin medications, and phentermine. 	E, Omega-3 fish oil, flax seed oil, and cod liver oil, herbal
3 DAYS BEFORE PROCEDURE:	
 Stop 325 mg aspirin and other NS. Celebrex, Mobic etc.), unless specification. Confirm your ride from procedure to 	
If you have diabetes, oral hypoglycemics may need	d to be held in the morning of the procedure. Please check with the
physician who prescribed this medication. Insulin ma	
THE DAY BEFORE YOUR PROCEDURE:	*NO SOLID FOODS*
When you wake up: You may only cor	nsume <u>CLEAR LIQUIDS all day</u> . Blue, white, yellow and green flavors
only. Water, sparkling/ vitamin water, sports drir cream/milk), soda, Jell-O, popsicles. Stay wel l	nk, apple/ white grape juice, broth/ bouillon, tea or coffee (without I hydrated.
STEP 1: At 5:00 pm - Open 1 bottle of	of 12 tablets.
	h 16 oz. of water (to the fill line). Swallow 1 tablet every 5 minutes ourse of one hour. Drink all 16 oz of water.
STEP 3: Between 6:30-7:30 pm - Drin	ok 2 additional 16 oz. glasses of water to dissolve the tablets you stay hydrated and keep drinking plenty of additional clear liquids
	ve steps, take the tablets and water slower. Peppermints may help to
SECOND DOSE: DATE	@
STEP 1: Repeat Step 1 and Step 2 from	n above.

STEP 2: Drink 2 additional 16 oz glasses of water to dissolve the tablets you have taken. It is important to stay hydrated and keep drinking plenty of additional clear liquids before bedtime.

If you become uncomfortable during any of the above steps, take the tablets and water slower. Peppermints may help to settle your stomach.

*If you take blood pressure or anxiety medications, take these with a small sip of water in the morning.

Do not chew gum, eat, or drink anything after completion of the prep, especially within 6 HOURS OF PROCEDURE

WHAT TO BRING/NOT TO BRING

Bring: Comfortable clothing; ID, insurance card, list of medications.

DO NOT bring: Jewelry, other valuables, contact lenses.

DO NOT apply lotions, creams, or ointments on your skin.

TRANSPORTATION

Please have a driver accompany you to be confirmed at check-in.

- *Without a confirmed driver during check-in, the procedure will not be able to be performed.
- *YOU WILL NOT BE ALLOWED TO DRIVE AFTER ANY PROCEDURE THAT REQUIRES SEDATION.
- *You must not operate a vehicle or heavy machinery, work, or drink alcohol for 12 hours after the procedure.
- *Someone should stay with you for 12 hours after the procedure.
- *You may resume regular activities, including driving, the next day.

IF YOU DO NOT SPEAK ENGLISH, PLEASE HAVE A TRANSLATOR ACCOMPANY YOU TO THE VISIT

CANCELLATION POLICY:

YOU WILL BE CHARGED A \$200.00 FEE IF YOU DO NOT CANCEL AT LEAST FIVE (5) BUSINESS DAYS PRIOR TO YOUR PROCEDURE.

BILLING/INSURANCE

- *All referrals (if needed) should be provided at least one week before your scheduled procedure.
- *Please notify us at least one week in advance of any insurance changes.
- *Please note that you may receive separate bills from pathology, anesthesiology, hospital facility, in addition to physician professional fees.

ANESTHESIA IS BILLED BY A THIRD PARTY – GREATER WASHINGTON ANESTHESIA ASSOCIATES Insurance may be out of network with anesthesia.

If you have any questions regarding your charges, please call 888-717-5383.

AT THE ENDOSCOPY UNIT

You will be escorted to a pre-procedure bay where you will change into a procedure gown. An IV will be placed. Your accompanying person/driver may wait with you until the procedure starts. During the procedure, they may wait in the waiting area or in the parking lot until the procedure is completed. After your procedure when you are awake and alert, your physician will review the findings with you. When you are ready to depart, you will be escorted out in a wheelchair.

Unexpected delays may occur during the procedure day due to medical needs of other patients. We will do our best to maintain scheduled times, but please be prepared for delays.

COLONOSCOPY CATEGORIES:

SCREENING COLONOSCOPY: Age 45 without any symptoms. No history of colon polyps, colorectal cancer, gastrointestinal symptoms, family history of advanced/ early colon polyps or colorectal cancer. If there is a family history of parents, siblings or children with colorectal cancer, a screening is recommended 10 years prior to the age of diagnosis of the family member. **SURVEILLANCE COLONOSCOPY:** Patient without symptoms but may have a personal history of GI disease such as ulcerative colitis or Crohn's Disease, colon polyps and/or colorectal cancer. Patients in this category are requested to undergo colonoscopy at shortened intervals. **Diagnostic testing copays and coinsurances may apply.**

DIAGNOSTIC COLONOSCOPY: Patient has current gastrointestinal symptoms, GI disease, iron deficiency anemia and/or any other abnormal tests or studies. **Diagnostic testing copays and coinsurances may apply.**

GASTROENTEROLOGY ASSOCIATES, PC www.doctorgi.com